

**Community Unit District 308**  
**Seventh Grade Honors Math Scope & Sequence**

- ◆ Standards taught and Assessed through Formative Assessments
- Standards taught and Assessed through Summative Assessments

[illegible]

**Community Unit District 308  
Seventh Grade Honors Math Scope & Sequence**

- ◆ Standards taught and Assessed through Formative Assessments  
● Standards taught and Assessed through Summative Assessments

Unit	1	2	3	4	5	6	7	8	9	10
<b>7.NS.A.3</b> Solve real-world and mathematical problems involving the four operations with rational numbers.		◆								
<b>The Number System</b>										
<b>8.NS.A Know that there are numbers that are not rational, and approximate them by rational numbers.</b>										
<b>8.NS.A.1</b> Know that numbers that are not rational are called irrational. Understand informally that every number has a decimal expansion; for rational numbers show that the decimal expansion repeats eventually, and convert a decimal expansion which repeats eventually into a rational number.		●								
<b>8.NS.A.2</b> Use rational approximations of irrational numbers to compare the size of irrational numbers, locate them approximately on a number line diagram, and estimate the value of expressions (e.g., $\pi^2$ ). For example, by truncating the decimal expansion of $\sqrt{2}$ , show that $\sqrt{2}$ is between 1 and 2, then between 1.4 and 1.5, and explain how to continue on to get better approximations.		◆	◆							
<b>Expressions &amp; Equations</b>										
<b>7.EE.A Use properties of operations to generate equivalent expressions.</b>										
<b>7.EE.A.1</b> Apply properties of operations as strategies to add, subtract, factor, and expand linear expressions with rational coefficients.										
<b>7.EE.A.2</b> Understand that rewriting an expression in different forms in a problem context can shed light on the problem and how the quantities in it are related. For example, $a + 0.05a = 1.05a$ means that "increase by 5%" is the same as "multiply by 1.05."										
<b>7.EE.B Solve real-life and mathematical problems using numerical and algebraic expressions and equations.</b>										
<b>7.EE.B.3</b> Solve multi-step real-life and mathematical problems posed with positive and negative rational numbers in any form (whole numbers, fractions, and decimals), using tools strategically. Apply properties of operations to calculate with numbers in any form; convert between forms as appropriate; and assess the reasonableness of answers using mental computation and estimation strategies. For example: If a woman making \$25 an hour gets a 10% raise, she will make an additional 1/10 of her salary an hour, or \$2.50, for a new salary of \$27.50. If you want to place a towel bar 9 3/4 inches long in the center of a door that is 27 1/2 inches wide, you will need to place the bar about 9 inches from each edge; this estimate can be used as a check on the exact computation.	◆									
<b>7.EE.B.4a</b> Use variables to represent quantities in a real-world or mathematical problem, and construct simple equations and inequalities to solve problems by reasoning about the quantities. <b>a.</b> Solve word problems leading to equations of the form $px + q = r$ and $p(x + q) = r$ , where $p$ , $q$ , and $r$ are specific rational numbers. Solve equations of these forms fluently. Compare an algebraic solution to an arithmetic solution, identifying the sequence of the operations used in each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?	◆									
<b>7.EE.B.4b</b> Use variables to represent quantities in a real-world or mathematical problem, and construct simple equations and inequalities to solve problems by reasoning about the quantities. <b>b.</b> Solve word problems leading to inequalities of the form $px + q > r$ or $px + q < r$ , where $p$ , $q$ , and $r$ are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.	◆									
<b>Expressions &amp; Equations</b>										
<b>8.EE.A Expressions and equations work with radicals and integer exponents.</b>										
<b>8.EE.A.1</b> Know and apply the properties of integer exponents to generate equivalent numerical expressions. For example, $3^2 \times 3^{-5} = 3^{-3} = 1/3^3 = 1/27$ .		●	●							
<b>8.EE.A.2</b> Use square root and cube root symbols to represent solutions to equations of the form $x^2 = p$ and $x^3 = p$ , where $p$ is a positive rational number. Evaluate square roots of small perfect squares and cube roots of small perfect cubes. Know that $\sqrt{2}$ is irrational.		●	●							
<b>8.EE.A.3</b> Use numbers expressed in the form of a single digit times an integer power of 10 to estimate very large or very small quantities, and to express how many times as much one is than the other. For example, estimate the population of the United States as 3 times $10^8$ and the population of the world as 7 times $10^9$ , and determine that the world population is more than 20 times larger.			●							
<b>8.EE.A.4</b> Perform operations with numbers expressed in scientific notation, including problems where both decimal and scientific notation are used. Use scientific notation and choose units of appropriate size for measurements of very large or very small quantities (e.g., use millimeters per year for seafloor spreading). Interpret scientific notation that has been generated by technology			◆							

**Community Unit District 308**  
**Seventh Grade Honors Math Scope & Sequence**

- ◆ Standards taught and Assessed through Formative Assessments
- Standards taught and Assessed through Summative Assessments

[illegible]

**Community Unit District 308**  
**Seventh Grade Honors Math Scope & Sequence**

- ◆ Standards taught and Assessed through Formative Assessments
- Standards taught and Assessed through Summative Assessments

	Unit	1	2	3	4	5	6	7	8	9	10
<b>8.G.A.4</b>	Understand that a two-dimensional figure is similar to another if the second can be obtained from the first by a sequence of rotations, reflections, translations, and dilations; given two similar two-dimensional figures, describe a sequence that exhibits the similarity between them.								●		
<b>8.G.A.5</b>	Use informal arguments to establish facts about the angle sum and exterior angle of triangles, about the angles created when parallel lines are cut by a transversal, and the angle-angle criterion for similarity of triangles. <i>For example, arrange three copies of the same triangle so that the sum of the three angles appears to form a line, and give an argument in terms of transversals why this is so.</i>				●						
<b>8.G.B Understand and apply the Pythagorean theorem.</b>											
<b>8.G.B.6</b>	Explain a proof of the Pythagorean Theorem and its converse.				◆						
<b>8.G.B.7</b>	Apply the Pythagorean Theorem to determine unknown side lengths in right triangles in real-world and mathematical problems in two and three dimensions.				●						
<b>8.G.B.8</b>	Apply the Pythagorean Theorem to find the distance between two points in a coordinate system.				●						
<b>8.G.C Solve real-world and mathematical problems involving volume of cylinders, cones, and spheres.</b>											
<b>8.G.C.9</b>	Know the formulas for the volumes of cones, cylinders, and spheres and use them to solve real-world and mathematical problems.		●								
<b>Statistics &amp; Probability</b>											
<b>7.SP.A Use random sampling to draw inferences about a population.</b>											
<b>7.SPA.1</b>	Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.										●
<b>7.SPA.2</b>	Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.										●
<b>7.SP.B Draw informal comparative inferences about two populations.</b>											
<b>7.SP.B.3</b>	Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability (mean absolute deviation) on either team; on a dot plot, the separation between the two distributions of heights is noticeable.										●
<b>7.SP.B.4</b>	Use measures of center and measures of variability for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.										●
<b>7.SP.C Draw informal comparative inferences about two populations.</b>											
<b>7.SP.C.5</b>	Understand that the probability of a chance event is a number between 0 and 1 that expresses the likelihood of the event occurring. Larger numbers indicate greater likelihood. A probability near 0 indicates an unlikely event, a probability around 1/2 indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event.										●
<b>7.SP.C.6</b>	Approximate the probability of a chance event by collecting data on the chance process that produces it and observing its long-run relative frequency, and predict the approximate relative frequency given the probability. For example, when rolling a number cube 600 times, predict that a 3 or 6 would be rolled roughly 200 times, but probably not exactly 200 times.										●
<b>7.SP.C.7</b>	Develop a probability model and use it to find probabilities of events. Compare probabilities from a model to observed frequencies; if the agreement is not good, explain possible sources of the discrepancy. <b>a.</b> Develop a uniform probability model by assigning equal probability to all outcomes, and use the model to determine probabilities of events. For example, if a student is selected at random from a class, find the probability that Jane will be selected and the probability that a girl will be selected.										●
<b>7.SP.C.7</b>	Develop a probability model and use it to find probabilities of events. Compare probabilities from a model to observed frequencies; if the agreement is not good, explain possible sources of the discrepancy. <b>b.</b> Develop a probability model (which may not be uniform) by observing frequencies in data generated from a chance process. For example, find the approximate probability that a spinning penny will land heads up or that a tossed paper cup will land open-end down. Do the outcomes for the spinning penny appear to be equally likely based on the observed frequencies?										●

**Community Unit District 308**  
**Seventh Grade Honors Math Scope & Sequence**

- ◆ Standards taught and Assessed through Formative Assessments
- Standards taught and Assessed through Summative Assessments

Unit	1	2	3	4	5	6	7	8	9	10
<b>7.SP.C.8</b> Find probabilities of compound events using organized lists, tables, tree diagrams, and simulation. <b>a.</b> Understand that, just as with simple events, the probability of a compound event is the fraction of outcomes in the sample space for which the compound event occurs.										●
<b>7.SP.C.8</b> Find probabilities of compound events using organized lists, tables, tree diagrams, and simulation. <b>b.</b> Represent sample spaces for compound events using methods such as organized lists, tables and tree diagrams. For an event described in everyday language (e.g., "rolling double sixes"), identify the outcomes in the sample space which compose the event.										●
<b>7.SP.C.8</b> Find probabilities of compound events using organized lists, tables, tree diagrams, and simulation. <b>c.</b> Design and use a simulation to generate frequencies for compound events. . For example, use random digits as a simulation tool to approximate the answer to the question: If 40% of donors have type A blood, what is the probability that it will take at least 4 donors to find one with type A blood?										●
<b>Statistics and Probability</b>										
<b>8.SP.A Investigate patterns of association in bivariate data.</b>										
<b>8.SP.A.1</b> Construct and interpret scatter plots for bivariate measurement data to investigate patterns of association between two quantities. Describe patterns such as clustering, outliers, positive or negative association, linear association, and nonlinear association.						●				
<b>8.SP.A.2</b> Know that straight lines are widely used to model relationships between two quantitative variables. For scatter plots that suggest a linear association, informally fit a straight line, and informally assess the model fit by judging the closeness of the data points to the line.						●				
<b>8.SP.A.3</b> Use the equation of a linear model to solve problems in the context of bivariate measurement data, interpreting the slope and intercept. <i>For example, in a linear model for a biology experiment, interpret a slope of 1.5 cm/hr as meaning that an additional hour of sunlight each day is associated with an additional 1.5 cm in mature plant height.</i>						●				
<b>8.SP.A.4</b> Understand that patterns of association can also be seen in bivariate categorical data by displaying frequencies and relative frequencies in a two-way table. Construct and interpret a two-way table summarizing data on two categorical variables collected from the same subjects. Use relative frequencies calculated for rows or columns to describe possible association between the two variables. <i>For example, collect data from students in your class on whether or not they have a curfew on school nights and whether or not they have assigned chores at home. Is there evidence that those who have a curfew also tend to have chores?</i>									●	